

2Pass4sure

2Pass4sure

HOME

ALL VENDORS

GUARANTEE

FAQ

TESTIMONIALS

CART (0)

Reliable Certification Exam Questions and Exam Dumps!

Everything you need to prepare, learn & pass your certification exam easily.

365 days free updates. First attempt guaranteed success.

Select a vendor...

Select an test...

Your email address

Free Download Demo

We're not the only ones **happy** about 2Pass4sure Practice Material ...

62819+ customers in 100+ countries use 2Pass4sure Test Engine. Meet our customers.

VOREED

GetCustom

JET ORANGE

iCompany

Paradoxx

iMessenger



<http://www.2pass4sure.com/>

Reliable Certification Exam Questions and Exam Dumps - 2Pass4sure

Exam : C2020-011

Title : IBM SPSS Statistics Level 1 v2

Vendors : IBM

Version : DEMO

NO.1 With IBM SPSS Statistics you can assign one of three levels of measurement to a variable. Which

is the correct list of all three levels?

- A. Nominal - Ordinal - Interval
- B. Categorical - Interval - Ratio
- C. Nominal - Ordinal - Scale
- D. Nominal - Interval – Scale

Answer: C

NO.2 If you request an aggregated dataset or data file in the Aggregate procedure, the number of cases

in the new aggregated file is equal to what?

- A. Number of cases in the original data file
- B. Number of aggregated summary variables
- C. Number of categories of the variables specified in the Break Variables list
- D. Number of cases that you specified in the Aggregate Data dialog box

Answer: C

NO.3 In the Aggregate data procedure¹ if you specify more than one Break Variable, _____ in the

Break Variables list will determine case order.

- A. Variable size
- B. Variable label
- C. Variable name
- D. Variable order

Answer: D

NO.4 If you use the IBM SPSS Statistics Select Cases dialog to conduct analysis on a subset of cases and you want to keep all cases in the same dataset, which option is appropriate?

- A. Copy unselected cases to a new dataset
- B. Copy selected cases to a new dataset
- C. Delete unselected cases
- D. Filter out unselected cases

Answer: D

NO.5 What type of graph would you use to see the distribution of a scale variable?

- A. Bar chart
- B. Pie chart
- C. Histogram
- D. Scatter plot

Answer: C

NO.6 In the output of the CROSSTABS procedure, large residuals indicate agreement between the data

and the null hypothesis of independence.

- A. True
- B. False

Answer: B

NO.7 When you read data stored in an Excel file via File ... Open ... Data, you may need to specify which options? (Choose three.)

- A. Read variable names from the first row of data
- B. Read a Range of cells
- C. Specify the Worksheet to read/import
- D. Number of Worksheets to read/import

Answer: A,B,C